

A note on Pollard's Theorem

Y. O. Hamidoune*

O. Serra†

Abstract

Let A, B be nonempty subsets of an abelian group G . Let $N_i(A, B)$ denote the set of elements of G having i distinct decompositions as a product of an element of A and an element of B . We prove that

$$\sum_{1 \leq i \leq t} |N_i(A, B)| \geq t(|A| + |B| - t - \alpha + 1 + w) - w,$$

where α is the largest size of a coset contained in AB and $w = \min(\alpha - 1, 1)$, with a strict inequality if $\alpha \geq 3$ and $t \geq 2$, or if $\alpha \geq 2$ and $t = 2$.

This result is a local extension of results by Pollard and Green–Ruzsa and extends also for $t > 2$ a recent result of Gryniewicz, conjectured by Dicks–Ivanov (for non necessarily abelian groups) in connection to the famous Hanna Neumann problem in Group Theory.

MSC Classification: 11B60, 11B34, 20D60.

1 Introduction

Groups will be written multiplicatively.

Let G be a group and let A, B be two nonempty subsets of G . The *representation function* $r_{A,B} : G \rightarrow \mathbb{N}$ is defined by the relation

$$r_{A,B}(x) = |(xB^{-1}) \cap A|.$$

The subgroup generated by a subset S will be denoted by $\langle S \rangle$.

Let X be a subset of an abelian group G . Recall that the subgroup $\{x \in G : Xx = X\}$ is called the *period* of X .

Our notations follow almost everywhere the terminology of Nathanson [10]. In particular we write

$$N_t(A, B) = \{x \in AB : r_{A,B}(x) \geq t\}.$$

Pollard [14] proved the following remarkable result:

*UPMC Univ Paris 06, E. Combinatoire, Case 189, 4 Place Jussieu, 75005 Paris, France.

†Universitat Politècnica de Catalunya, Jordi Girona, 1, E-08034 Barcelona, Spain.

Theorem A (Pollard [14]) *Let A and B be nonempty subsets of a group with a prime order p and let $t \leq \min(|A|, |B|)$. Then*

$$\sum_{1 \leq i \leq t} |N_i(A, B)| \geq t \min(p, |A| + |B| - t).$$

Very recently Nazarewicz, O'Brien, O'Neill and C. Staples [12] obtained the equality cases in Pollard's Theorem A. A restricted-sum version of the inequalities is obtained by Caldeira and Dias da Silva [1].

Under a Chowla's type condition, Pollard [15] proved a generalization of his theorem to composite moduli. He also suggested the problem of finding conditions implying the validity of his inequalities in an abelian group.

In connection with old questions on sum-free sets [5], the following generalization of Pollard's Theorem to arbitrary abelian groups was given by Green–Ruzsa:

Theorem B (Green–Ruzsa [5]) *Let G be an abelian group and let A and B be nonempty subsets of G and let $t \leq \min(|A|, |B|)$. Then*

$$\sum_{1 \leq i \leq t} |N_i(A, B)| \geq t \min(|G|, |A| + |B| - \mu(G) - t + 1),$$

where $\mu(G)$ denotes the cardinality of the largest subgroup of G different from G .

In the prime case $\mu(G) = 1$ and hence Green–Ruzsa Theorem reduces to Pollard's Theorem A.

Kemperman [7] proved a non-abelian counter-part of Kneser's Theorem (see Theorem C in Section 2) stating that there is a finite subgroup H such that $|AB| \geq |A| + |B| - |H|$, and $aHb \subset AB$ for some $a \in A$ and $b \in B$. Dicks–Ivanov [3] obtained a generalization of this result in connection with a famous problem in Group Theory and formulated the following:

Conjecture (Dicks–Ivanov [2, 4]) *Let A, B be finite nonempty subsets of a group G with $2 \leq |B| \leq |A|$. Then one of the following conditions holds:*

- (i) $|N_1(A, B)| + |N_2(A, B)| \geq 2(|A| + |B| - 2)$,
- (ii) $N_2(A, B)$ contains a left coset with cardinality ≥ 3 .

Gryniewicz [6, Theorem 3.1] proved that $|N_1(A, B)| + |N_2(A, B)| \geq 2(|A| + |B| - \max(2, |H|))$, where H is the period of $N_2(A, B)$. Clearly this result implies the validity of the Dicks–Ivanov Conjecture in the abelian case. Two easier proofs of this conjecture in the abelian case were proposed later by Dicks–Ivanov [4] and by the authors. Also Gryniewicz investigates in [6] conditions implying the validity of the inequality $\sum_{1 \leq i \leq t} |N_i(A, B)| \geq t(|A| + |B| - 2t + 1)$.

Let $\alpha(A, B)$ denote the cardinality of the largest left coset (non necessarily proper) contained in AB . In the spirit of the Dicks–Ivanov Conjecture, we prove the next local Green–Ruzsa type Theorem:

Theorem 1 *Let t be an integer and A, B be finite subsets of an abelian group G with $1 \in A \cap B$ and $|A| \geq |B| \geq t \geq 1$. Set $\alpha = \alpha(A, B)$ and $w = \min(\alpha - 1, 1)$. Then*

$$\sum_{1 \leq i \leq t} |N_i(A, B)| \geq t(|A| + |B| - t - \alpha + 1 + w) - w. \quad (1)$$

Moreover the inequality (1) is strict in the following cases:

(I) $\alpha \geq 3$ and $t \geq 2$,

(II) $\alpha \geq 2$ and $t = 2$.

For $\alpha = 1$ or $t = 1$, the inequality (1) is not necessarily strict. By Theorem 1, the inequality (1) is strict for $t > 1$, unless $t \geq 3$ and $\alpha = 2$. We suspect that it is still strict in this case as well.

For $t = 2$ Theorem 1, follows from the result of Gryniewicz mentioned above.

2 Preliminaries

We shall need the following results:

Lemma 2 (folklore) *Let G be a finite abelian group. Let A, B be nonempty subsets such that $|A| + |B| \geq |G| + t$. Then for every $x \in G$, $r_{A,B}(x) \geq t$. In particular $\sum_{x \in G} r_{A,B}(x) \geq t|G|$.*

Proof. We have clearly $|(xB^{-1}) \cap A| \geq t$. $\square$

Theorem C (Kneser [9, 10]) *Let G be an abelian group and let $A, B \subset G$ be finite subsets of G with $|AB| \leq |A| + |B| - 2$. Let H be the period of AB . Then $|H| \geq 2$. Moreover*

$$|AB| = |AH| + |BH| - |H|, \quad (2)$$

$$N_2(A, B) = AB. \quad (3)$$

The equality (3) is known as the Kemperman-Scherk Theorem. It is proved by Scherck [13] in the abelian case and by Kemperman [7] in the non-abelian one. But in the abelian case it can be easily derived from (2) as follows:

Take $x \in AB$. Then $x \in A_1 B_1$, where A_1 (resp. B_1) is the trace of some coset on A (resp. B). Now $2|H| - |A_1| - |B_1| \leq |AH| + |BH| - |A| - |B| = |AB| + |H| - |A| - |B| \leq |H| - 2$. It follows that $|xA^{-1} \cap B| \geq 2$. $\square$

The following lemma is implicitly proved by Pollard in [14]. We give few hints for its proof in order to make the present work self-contained:

Lemma 3 *Let G be an abelian group. Let A, B be nonempty sets with $|B| \leq |A|$. Let t, v be integers such that $0 \leq v \leq t$ and let $z \in G$.*

$$|N_t(A, B)| = |N_t(zA, B)|, \quad (4)$$

$$\sum_{1 \leq i \leq |B|} |N_i(A, B)| = |A||B|, \quad (5)$$

$$\sum_{1 \leq i \leq t} |N_i(A, B)| \geq \sum_{1 \leq i \leq v} |N_i(A \cap B, A \cup B)| + \sum_{1 \leq i \leq t-v} |N_i(A \setminus B, B \setminus A)|. \quad (6)$$

Proof. The proof of (i) is obvious. We have

$$\sum_{1 \leq i \leq |B|} |N_i(A, B)| = \sum_{x \in G} r_{A, B}(x) = \sum_{x \in A} |xB| = |A||B|,$$

and hence (ii) holds. We have also

$$\begin{aligned} \sum_{1 \leq i \leq t} |N_i(A, B)| &= \sum_{x \in G} \min(t, r_{A, B}(x)) \\ &= \sum_{x \in G} \min(t, r_{A \setminus B, B \setminus A}(x) + r_{A \cap B, B \setminus A}(x) + r_{A \cap B, B \cap A}(x) + r_{A \setminus B, A \cap B}(x)) \\ &= \min(t, r_{A \setminus B, B \setminus A}(x) + r_{A \cap B, A \cup B}(x)) \\ &\geq \sum_{1 \leq i \leq v} |N_i(A \cap B, A \cup B)| + \sum_{1 \leq i \leq t-v} |N_i(A \setminus B, B \setminus A)|. \end{aligned}$$

□

We need the following lemma:

Lemma 4 *Let A, B be finite nonempty subsets of an abelian group G and let $a \in G$. Then*

- (i) $N_t(aA, B) = aN_t(A, B)$,
- (ii) $N_t((A \cap B), (A \cup B)) \subset N_t(A, B)$.
- (iii) $\alpha(aA \cup B, aA \cap B) \leq \alpha(A, B)$.

The proof is easy.

3 Proof of Theorem 1

The proof is by induction on $|B|$.

If $|B| = t$ the result follows from (5) and the inequality is strict.

Assume first that $AB = A$. Then A has a decomposition $A = A_1 \cup \dots \cup A_k$ as the union of $\langle B \rangle$ -cosets, and $\alpha \geq |\langle B \rangle| \geq |B|$.

By Lemma 2 we have $\sum_{1 \leq i \leq t} N_i(A_j, B) = t|A_j|$ and

$$\sum_{1 \leq i \leq t} |N_i(A, B)| = \sum_{1 \leq i \leq t, 1 \leq j \leq k} |N_i(A_j, B)| = t|A|,$$

which proves the result and the inequality (1) is strict if $t > 1$. So we may assume $AB \neq A$.

Then there is $b \in B$ and $a \in A$ such that $b \notin Aa^{-1}$. In particular $1 \leq |B \cap (Aa^{-1})| < |B|$. Put $A' = Aa^{-1}$ and $v = |A' \cap B|$. Set $S = A' \setminus B$ and $T = B \setminus A'$.

Using Lemma 3 and Lemma 4, for every u with $0 \leq u \leq t$, we have

$$\sum_{1 \leq i \leq t} |N_i(A, B)| \geq \sum_{1 \leq i \leq u} |N_i(A' \cap B, A' \cup B)| + \sum_{1 \leq i \leq t-u} |N_i(S, T)|. \quad (7)$$

Suppose first that $t \leq v$. Then, by (7) applied with $u = t$ and the induction hypothesis, we have

$$\sum_{1 \leq i \leq t} |N_i(A, B)| \geq t(|A| + |B| - t - \alpha + 1 + w) - w,$$

Also the inequality (1) is strict in case (II) by induction and in case (I) either by induction or if $2 = \alpha(A' \cup B, A' \cap B) < \alpha$.

Suppose now that $t > v$. Since $v = |B \cap A'|$, we have by (5)

$$\sum_{1 \leq i \leq v} |N_i(A' \cap B, A' \cup B)| = v|A' \cup B| = v(|A| + |B| - v). \quad (8)$$

We now estimate the second summand in the left hand side of (7) with $u = v$.

By the induction hypothesis we have

$$\sum_{1 \leq i \leq t-v} |N_i(S, T)| \geq (t-v)(|A| + |B| - 2v - (t-v) - \alpha + 1 + w) - w. \quad (9)$$

By adding the second terms in (8) and (9) we have,

$$\begin{aligned} \sum_{1 \leq i \leq t} |N_i(A, B)| &\geq v(|A| + |B| - v) + (t-v)(|A| + |B| - t - v - \alpha + 1 + w) - w \\ &= t(|A| + |B| - t - \alpha + 1 + w) + v(\alpha - w - 1) - w. \end{aligned}$$

The inequality (1) is clearly strict if $\alpha \geq 3$.

In order to complete the proof we have only to check that this inequality is strict for $t = \alpha = 2$ when $v = 1 < t$. Suppose this is not the case. From equality in (9) we have $|ST| = |S| + |T| - 2$. By Kneser Theorem, both S and T are H -periodic by a subgroup H of cardinality $|H| = \alpha = 2$.

It follows by (3) that

$$|N_2(A', B)| \geq |N_2(S, T)| = |N_1(S, T)| = |S| + |T| - 2 = |A| + |B| - 4.$$

Moreover we may assume $|N_1(A, B)| \leq |A| + |B| - 1$ since otherwise $|N_1(A, B)| + |N_2(A, B)| \geq 2(|A| + |B| - 2)$ and we are done.

Hence $|N_1(A', B) \setminus ST| = 3$, and since S and T are H -periodic, $N_1(A', B) \setminus ST = \{1\} \cup aH$, for some a . We shall consider only the case $a \in S$, the other case being essentially the same.

Observe that $aH \cap A'T = \emptyset$, otherwise $aH \subset N_2(A', B)$ because $aH = aH.1$. Also $1 \notin A'T = A'TH$. It follows that $|A'T| \leq |A'B| - 3$, and hence $A'T = ST$. By Kneser's Theorem C,

$$|ST| = |A'T| \geq |A'H| + |HT| - |H| = |A| + |H| + |T| - |H| = |S| + |T| - 1,$$

a contradiction. This completes the proof. $\square$

Acknowledgement

The authors would like to thank Professor W. Dicks for stimulating discussions and comments on Pollard's inequality and also for several remarks allowing us to remove some obscure parts in our first draft.

References

- [1] C. Caldeira and J. A. Dias da Silva, A Pollard type result for restricted sums, *J. Number Theory* 72 (1998), no. 2, 153-173.
- [2] W. Dicks, Oral communications.
- [3] W. Dicks, S. V. Ivanov, On the intersection of free subgroups in free products of groups, *Math. Proc. Cambridge Philos. Soc.* (to appear). <http://mat.uab.cat/~dicks/ivanov.html>
<http://arxiv.org/abs/math.GR/0702363>.
- [4] W. Dicks, The abelian case of a combinatorial conjecture of Dicks and Ivanov, Notes for seminar March 27, 2008
- [5] B. Green, I.Z. Ruzsa, Sum-free sets in abelian groups. *Israel J. Math.* 147 (2005), 157–188.
- [6] David J. Grynkiewicz, Extending Pollard's Theorem for t-representable sums, <http://arxiv.org/abs/0803.2601> preprint.
- [7] J.H.B. Kempermann, On complexes in a semigroup, *Nederl. Akad. Wetensch. Proc. Ser. A.* 59 *Indag. Math.* 18 (1956), 247–254.
- [8] M. Kneser, Abschätzung der asymptotischen Dichte von Summenmengen. *Math. Z.* 58, (1953). 459–484.

- [9] M. Kneser, Summenmengen in lokalkompakten abelschen Gruppen. *Math. Zeit.* **66** (1956), 88–110.
- [10] M. B. Nathanson, *Additive Number Theory. Inverse problems and the geometry of sumsets*, Grad. Texts in Math. 165, Springer, 1996.
- [11] P. Scherk, J. H. B. Kemperman, Complexes in abelian groups. *Canadian J. Math.* **6**, (1954). 230–237.
- [12] E. Nazarewicz, M. O’Brien, M. O’Neill and C. Staples, Equality in Pollard’s theorem on set addition of congruence classes. *Acta Arith.* to appear.
- [13] P. Scherk, L. Moser, Advanced Problems and Solutions: Solutions: 4466, *Amer. Math. Monthly* **62** (1955), no. 1, 46–47.
- [14] J. M. Pollard, A generalisation of the theorem of Cauchy and Davenport, *J. London Math. Soc.* (2) **8** (1974), 460–462.
- [15] J. M. Pollard, Addition properties of residue classes, *J. London Math. Soc.* (2) **11** (1975), no. 2, 147–152.